

Ovarian Cancer Red Flags



Be aware!

**New diagnosis
of IBS in the >50s**



Be aware!

**New onset 'overactive bladder'/
'recurrent UTI' in >50s**

- ▶ Ovarian cancer may be misdiagnosed as irritable bowel syndrome (IBS), overactive bladder or recurrent urinary tract infection.
- ▶ First presentation with IBS in the >50s is rare.
- ▶ Women in this age group who have experienced recurrent symptoms within the last 12 months that suggest IBS should be offered a CA125 blood test and considered for USS pelvis and abdomen.
- ▶ Recurrent symptoms of 'urinary tract infection' should always be investigated by sending the mid-stream urine for culture.
- ▶ If the urine is repeatedly sterile, and the woman is >50, a CA125 blood test and USS pelvis and abdomen should be considered.



**Target
Ovarian
Cancer**

targetovariancancer.org.uk

Update your knowledge of ovarian cancer. Take our free accredited training courses, download fact sheets, watch and listen to our expert videos and podcasts and join our GP network for the latest in ovarian cancer education.

targetovariancancer.org.uk/health-professionals

Authors: Dr Charlotte Bădescu and Dr Eloise Elphinstone



30 Angel Gate, London, EC1V 2PT



020 7923 5470



info@targetovariancancer.org.uk



TargetOvarianCancer



@TargetOvarian



@TargetOvarian



targetovariancancer.org.uk